

## Imaging Study Sheds New Light on Alcohol-Related Birth Defects

---



A collaborative research effort by scientists at the University of North Carolina School of Medicine, Duke University, and University College of London in the UK, sheds new light on alcohol-related birth defects.

The project, led by Kathleen K. Sulik, PhD, a professor in the Department of Cell and Developmental Biology and the Bowles Center for Alcohol Studies at UNC, could help enhance how doctors diagnose birth defects caused by alcohol exposure in the womb. The findings also illustrate how the precise timing of that exposure could determine the specific kinds of defects.

Fetal alcohol syndrome (FAS) is at the severe end of fetal alcohol spectrum disorders (FASD). First described in 1972, FAS is recognized by a specific pattern of facial features: small eyelid openings, a smooth ridge on the upper lip (absence of a central groove, or philtrum), and a thin upper lip border.

In its full-blown state, FAS affects roughly 1 in 750 live births in the U.S. And while clinicians typically look for those classical facial features in making a diagnosis, within the broader classification of FASD "adverse outcomes vary considerably and most individuals don't exhibit the facial characteristics that currently define FAS," said the study's lead author Robert J. Lipinski, PhD, a postdoctoral scientist in Sulik's lab. "This study could expand the base of diagnostic criteria used by clinicians who suspect problems caused by maternal alcohol use."

In the study, described in the August 22, 2012 issue of the online journal PLOS ONE, Lipinski and Sulik treated one group of mice with alcohol on their seventh day of pregnancy, a time corresponding to the third week of pregnancy in humans. A second group of mice was treated just 36 hours later, approximating the fourth week of human pregnancy. The amount of alcohol given was large, "high doses that most women wouldn't achieve unless they were alcoholic and had a tolerance for alcohol," Sulik said.

Near the end of pregnancy, the fetuses were then imaged at Duke University. These 3D data sets showed individual brain regions, as well as accurate and detailed facial surfaces, from which Hammond and research assistant and co-author Michael Suttie performed shape analyses.

## Imaging Study Sheds New Light on Alcohol-Related Birth Defects

---

The team found that the earlier alcohol exposure time elicited the classic FAS facial features, including characteristic abnormalities of the upper lip and eyes. What they observed in fetuses exposed just 36 hours later, however, was a surprise. These mice exhibited unique and in some cases opposing facial patterns, such as shortened upper lip, a present philtrum, and the brain, instead of appearing too narrow in the front, appeared wide.

The project was supported by the National Institute of Alcoholism and Alcohol Abuse, the National Institute of Biomedical Imaging and Bioengineering and was conducted in conjunction with the Collaborative Initiative on Fetal Alcohol Spectrum Disorders (CIFASD).

To learn more about Alcohol and Pregnancy - Fetal Alcohol Effects (FAE), [click here](#).

*Citation: Lipinski RJ, Hammond P, O'Leary-Moore SK, Ament JJ, Pecevich SJ, et al. (2012) Ethanol-Induced Face-Brain Dysmorphology Patterns Are Correlative and Exposure-Stage Dependent. PLoS ONE 7(8): e43067. doi:10.1371/journal.pone.0043067*

*Source: University of North Carolina School of Medicine*